

WHAT ARE CO-OCCURRING CONDITIONS ('COMORBIDITY')?

PAGE SUMMARY




- On this website, 'comorbidity' refers to the occurrence of a substance use disorder and a mental health disorder, or symptoms of these disorders, at the same time. Another term used for this is 'co-occurring substance use and mental health disorders'.
- There are several reasons why substance use and mental health disorders might occur together. Once established, they often interact in ways that maintain one another, making recovery difficult if treatment only focuses on one of these conditions.
- People with co-occurring substance use (e.g. [crystal methamphetamine 'ice' dependence](#)) and mental health disorders generally experience poorer overall health and wellbeing compared to those who have just one of these conditions.
- It is important that treatment services attend to both substance use *and* mental health problems when discussing treatment with their clients.



WHAT ARE CO-OCCURRING CONDITIONS ('COMORBIDITY')?

'Comorbidity' occurs when a person experiences two or more medical conditions at the same time. These co-occurring conditions are referred to as 'comorbid' and often interact with one another over long periods of time. There are many different types of comorbidity that people can experience.

Here we focus on the type of comorbidity where someone experiences both a substance use disorder (e.g. [crystal methamphetamine 'ice' dependence](#)) and a mental health disorder (e.g. **anxiety, depression, psychosis**), or symptoms of these disorders, at the same time.



Get the Factsheet: What are co-occurring conditions ('comorbidity')?


WATCH VIDEO: WHAT IS COMORBIDITY?



This video was funded by the [Australian Government Department of Health](#). The video was delivered by researchers from the [Matilda Centre for Research in Mental Health and Substance Use at the University of Sydney](#).

HOW DO CO-OCCURRING CONDITIONS DEVELOP?

There are several possible reasons why two or more disorders may occur at the same time.




One condition may directly cause the other. Someone may become **dependent** on alcohol and/or other drugs (like ice) because they use them to help cope with mental health symptoms. This is often referred to as “self-medication”. Alternatively, someone’s use of alcohol and/or other drugs may impair the way their brain functions, leading to poor mental health. For example, **withdrawal** from ice can leave people feeling depressed due to changes in brain chemicals, and heavy, consistent use of ice can sometimes cause acute **psychotic** reactions. For

most people these symptoms may subside after a few hours or a few days, however a small number of people may find these symptoms last much longer (e.g. more than a few weeks) or continue even when they are no longer using ice. This might mean that an underlying psychotic disorder, such as **schizophrenia**, is present.

One condition may indirectly cause the other. Poor mental health may cause life difficulties that lead someone to use alcohol and/or other drugs. For example, the experience of mental health problems may limit someone's ability to study or work. Someone in this position may start using alcohol or drugs to manage the stress of not being able to study or work how they would like to. In the opposite direction, using alcohol and/or other drugs may limit someone's ability to study or work. The stress of not being able to study or work how they would like to, may then impair their mental health.

Both conditions may be caused by something else. Sometimes two conditions can be caused by a shared biological, psychological, social or environmental risk factor. A shared risk factor is something about a person or their circumstances that increases their risk of experiencing each of the conditions. For example, both substance use disorders and mental health conditions have been associated with cognitive impairment and lower socioeconomic status.



A lot of research has been conducted to determine which disorder typically develops first out of the different co-occurring substance use and mental health disorders people experience. The evidence so far is not consistent, and researchers have found differences depending on a person's gender as well as the specific types of mental health disorder/s people are experiencing.

Understanding when different substance use and mental health disorders develop and how they interact with one another can help us to understand how best to prevent them from occurring. But sometimes this is tricky to work this out – it can be a bit like the 'chicken or the egg' example when trying to work out if a mental health disorder has come first, or whether a substance use disorder has come first. This can also be tricky because often the symptoms of mental and substance use disorders overlap and emerge at similar points in time during adolescence.


Regardless of which disorder has come first, once these disorders have developed, they most likely maintain and exacerbate one another. This can make it difficult for people to recover if they aren't able to access treatment for both conditions. For example, someone may use alcohol and/or other drugs to reduce symptoms of **anxiety**, yet research shows that repeated use of the substances can lead to increased anxiety for people. Regardless of what order these different disorders develop in, the strategies used to manage them remain the same.

HOW COMMON ARE CO-OCCURRING CONDITIONS?

Substance use disorders and mental health disorders are both common in Australia. National population estimates indicate that one in two Australians will develop a substance use, **anxiety** or mood disorder (e.g. **depression**) in their lifetime, and one in five meet criteria for a substance use, anxiety or mood disorder annually.



The co-occurrence of substance use disorders and mental health disorders is common. Mental Health and Wellbeing approximately 1 in 3 individuals with a substance use disorder also have a mood disorder. Rates are even higher among people who are in alcohol and other drug treatment services, with Australian AOD treatment services having at least one co-occurring mental health disorder.



WHAT ARE THE HARMS ASSOCIATED WITH CO-OCCU

People with co-occurring substance use and mental health disorders generally experience poorer overall health and wellbeing compared to those experiencing single disorders. Co-occurring mental health conditions can also place an enormous strain on a person’s family and others close to them, both emotionally and financially. See the figure below for a summary of the harms associated with comorbidity.



Adapted from the Comorbidity Guidelines. See the key source list below for more information.

DO CO-OCCURRING CONDITIONS AFFECT TREATMENT?

Research into whether people who have a co-occurring mental health condition respond differently to alcohol and other drug treatment is mixed. Some research suggests people who enter alcohol and other drug (AOD) treatment with a co-occurring mental health disorder do not respond as well to treatment as those without these conditions. Other studies have shown people with a co-occurring mental health disorder benefit as much from AOD treatment as those who do not. Research also suggests, however, that although these people benefit just as much from AOD treatment, they continue to experience more serious substance use problems, as well as poorer physical and mental health and overall functioning compared to others when they exit treatment.

On balance, the evidence indicates that **having co-occurring conditions is not an insurmountable barrier to treating people with AOD use disorders**, however some clients with co-occurring mental health disorders may require additional treatment. Some interventions have been developed to specifically treat certain comorbidities, however more research is needed before they can be used in practice.



People experiencing co-occurring substance use and mental health issues should discuss this with their local doctor, AOD counsellor or mental health professional to ensure both areas of concern are addressed in treatment.

WHAT MENTAL HEALTH PROBLEMS CAN CO-OCCUR WITH ICE USE?

People who use ice may experience co-occurring mental health problems such as **anxiety**, **depression** and **psychosis**. For more information visit [mental health effects of ice](#). For more information about treatment and support services available visit our [get help page](#).

KEY SOURCE

Information on this webpage has been adapted from the Guidelines on the management of co-occurring alcohol and other drug and mental health conditions in alcohol and other drug treatment settings (2nd edition).

[Read the full Comorbidity Guidelines, link opens in a new window.](#)

See full reference list below.

REFERENCES

- Burns, L., Teesson, M., & O'Neill, K. (2005). The impact of comorbid anxiety and depression on alcohol treatment outcomes. *Addiction*, 100, 787-796. doi: [10.1111/j.1360-0443.2005.001069.x](https://doi.org/10.1111/j.1360-0443.2005.001069.x)

- Callaly, T., Trauer, T., Munro, L., & Whelan, G. (2001). Prevalence of psychiatric disorder in a methadone maintenance population. *Aust N Z J Psychiatry*, 35(5), 601-5. doi: [10.1080/0004867010060507](https://doi.org/10.1080/0004867010060507)
- Darke S., Kaye S., McKetin R., & Duflou J. (2008). Major physical and psychological harms of methamphetamine use. *Drug Alcohol Rev* 2008, 27, 253–62. doi: [10.1080/09595230801923702](https://doi.org/10.1080/09595230801923702)
- Dickey, B., Normand, S. L., Weiss, R. D., Drake, R. E., & Azeni, H. (2002). Medical morbidity, mental illness, and substance use disorders. *Psychiatric Services*, 53, 861-867. doi: [10.1176/appi.ps.53.7.861](https://doi.org/10.1176/appi.ps.53.7.861)
- Fiorentini, A., Sara Volonteri, L. S., Dragogna, F., Rovera, C., Maffini, M., Mauri, M.C., & Altamura, C.A. (2011). Substance-induced psychoses: A critical review of the literature. *Current Drug Abuse Reviews*, 4, 228-240. PMID: 21999698
- Fletcher, J. M. (2008). Adolescent depression: Diagnosis, treatment, and educational attainment. *Health Economics*, 17, 1215-1235. doi: [10.1002/hec.1319](https://doi.org/10.1002/hec.1319)
- Hildebrand, A., Behrendt, S., & Hoyer, J. (2015). Treatment outcome in substance use disorder patients with and without comorbid posttraumatic stress disorder: A systematic review. *Psychotherapy Research*, 25, 565-582. doi: [10.1080/10503307.2014.923125](https://doi.org/10.1080/10503307.2014.923125)
- Horwood, L. J., Fergusson, D. M., Hayatbakhsh, M. R., Najman, J. M., Coffey, C., Patton, G. C., . . . Hutchinson, D. M. (2010). Cannabis use and educational achievement: Findings from three Australasian cohort studies. *Drug and Alcohol Dependence*, 110, 247-253. doi: [10.1016/j.drugalcdep.2010.03.008](https://doi.org/10.1016/j.drugalcdep.2010.03.008)
- Johnson, M. E., Brems, C., & Burke, S. (2002). Recognizing comorbidity among drug users in treatment. *American Journal of Drug and Alcohol Abuse*, 28, 243-261.
- Kavanagh, D. J. & Connolly, J. M. (2009). Interventions for co-occurring addictive and other mental disorders (AMDs). *Addictive Behaviors*, 34, 838-845. doi: [10.1016/j.addbeh.2009.03.005](https://doi.org/10.1016/j.addbeh.2009.03.005)
- Kessler, R.C., Berglund, P., Demler, O., Jin, R., Merikangas, K. R., Walters, E. E. (2005). Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry*, 2005, 62(6), 593-602. doi: [10.1001/archpsyc.62.6.593](https://doi.org/10.1001/archpsyc.62.6.593)
- Kessler, R. C. (1995). Epidemiology of psychiatric comorbidity, in Textbook in psychiatric epidemiology, M.T. Tsuang, M. Tohen, and G.E.P. Zahner, Editors. New York, NY: Wiley and Sons, 179-197.
- Kessler, R. C., Foster, C. L., Saunders, W. B., & Stang, P. E. (1995). Social consequences of psychiatric disorders, I: Educational attainment. *American Journal of Psychiatry*, 152, 1026-1032. doi: [10.1176/ajp.152.7.1026](https://doi.org/10.1176/ajp.152.7.1026)
- Khantzian, E. J., & Albanese, M. J. (2008). Understanding addiction as self medication: Finding hope behind the pain. Lanham, M. D: Rowman and Littlefield Publishers.
- Kingston, R. E. F., Marel, C., & Mills, K. L. (2016). A systematic review of the prevalence of comorbid mental health disorders in people presenting for substance use treatment in Australia. *Drug and Alcohol Review*, 36, 527-539. doi: [10.1111/dar.12448](https://doi.org/10.1111/dar.12448)
- Leeies, M., Pagura, J., Sareen, J., & Bolton, J.M. (2010). The use of alcohol and drugs to self-medicate symptoms of posttraumatic stress disorder. *Depression and Anxiety*, 27, 731-736. doi: [10.1002/da.20677](https://doi.org/10.1002/da.20677)
- Lloyd, C. (1998). Risk factors for problem drug use: Identifying vulnerable groups. *Drugs: Education, Prevention and Policy*, 5, 217-232. doi: [10.3109/09687639809034084](https://doi.org/10.3109/09687639809034084)
- Lubman, D.I., Allen, N. B., Rogers, N., Cementon, E., & Bonomo, Y. (2007). The impact of co-occurring mood and anxiety disorders among substance-abusing youth. *Journal of Affective Disorders*, 2007, 103(1-3), 105-112. doi: [10.1016/j.jad.2007.01.011](https://doi.org/10.1016/j.jad.2007.01.011)

[10.1016/j.jad.2007.01.011](https://doi.org/10.1016/j.jad.2007.01.011)

- Marel, C., Mills, K.L., Kingston, R., Gournay, K., Deady, M., Kay-Lambkin, F., Baker, A., & Teesson, M. (2016). Guidelines on the management of co-occurring alcohol and other drug and mental health conditions in alcohol and other drug treatment settings (2nd edition). Sydney, Australia: Centre of Research Excellence in Mental Health and Substance Use, National Drug and Alcohol Research Centre, University of New South Wales (now Matilda Centre for Research in Mental Health and Substance Use, University of Sydney).
- Milby, J. B., Conti, K., Wallace, D., Mennemeyer, S., Mrug, S., & Schumacher, J. E. (2015). Comorbidity effects on cocaine dependence treatment and examination of reciprocal relationships between abstinence and depression. *Journal of Consulting and Clinical Psychology, 83*, 45-55. doi: [10.1037/a0037960](https://doi.org/10.1037/a0037960)
- Mills, K. L., Teesson, M., Ross, J., & Darke, S. (2007). The impact of post-traumatic stress disorder on treatment outcomes for heroin dependence. *Addiction, 102*, 447-454. doi: [10.1111/j.1360-0443.2006.01711.x](https://doi.org/10.1111/j.1360-0443.2006.01711.x)
- Ouimette, P., Goodwin, E., & Brown, P. J. (2006). Health and well being of substance use disorder patients with and without posttraumatic stress disorder. *Addictive Behaviors, 31*, 1415-1423. doi: [10.1016/j.addbeh.2005.11.010](https://doi.org/10.1016/j.addbeh.2005.11.010)
- Robinson, J., Sareen, J., Cox, B., & Bolton, J. (2011). Role of self-medication in the development of comorbid anxiety and substance use disorders: A longitudinal investigation. *Archives of General Psychiatry, 75*, 800-807. doi: [10.1001/archgenpsychiatry.2011.75](https://doi.org/10.1001/archgenpsychiatry.2011.75)
- Schäfer, I., & Najavits, L. M. (2007). Clinical challenges in the treatment of patients with posttraumatic stress disorder and substance abuse. *Current Opinion in Psychiatry, 20*, 614-618. doi: [10.1097/YCO.0b013e3282f0ffd9](https://doi.org/10.1097/YCO.0b013e3282f0ffd9)
- Siegfried, N. (1998). A review of comorbidity: Major mental illness and problematic substance use. *Australian and New Zealand Journal of Psychiatry, 32*, 707-717. doi: [10.3109/00048679809113127](https://doi.org/10.3109/00048679809113127)
- Slade, T., McEvoy, P. M., Chapman, C., Grove, R., & Teesson, M. (2015). Onset and temporal sequencing of lifetime anxiety, mood and substance use disorders in the general population. *Epidemiology and Psychiatric Sciences, 2015*, 24(1), 45-53. doi: [10.1017/S2045796013000577](https://doi.org/10.1017/S2045796013000577)
- Slade, T., Johnston, A., Teesson, M., Whiteford, H., Burgess, P., Pirkis, J., & Saw, S. (2009). The mental health of Australians 2: Report on the 2007 National Survey of Mental Health and Wellbeing. Canberra, Australia: Australian Government Department of Health and Ageing.
- Slade, T., Johnston, A., Oakley Browne, M. A., Andrews, G., & Whiteford, H. (2007). National Survey of Mental Health and Wellbeing: Methods and key findings. *Australian and New Zealand Journal of Psychiatry, 43*, 594-605. doi: [10.1080/00048670902970882](https://doi.org/10.1080/00048670902970882)
- Stockwell, T., Hodgson, R., & Rankin, H. (1982). Tension reduction and the effects of prolonged alcohol consumption. *British Journal of Addiction, 1982*, 77, 65-73. doi: [10.1111/j.1360-0443.1982.tb03250.x](https://doi.org/10.1111/j.1360-0443.1982.tb03250.x)
- Teesson, M., Slade, T., & Mills, K. (2009). Comorbidity in Australia: Findings of the 2007 National Survey of Mental Health and Wellbeing. *Australian and New Zealand Journal of Psychiatry, 43*, 606-614. doi: [10.1080/00048670902970908](https://doi.org/10.1080/00048670902970908)
- Teesson, M., Degenhardt, L., Proudfoot, H., Hall, W., & Lynskey, M. (2005). How common is comorbidity and why does it occur? *Australian Psychologist, 40*, 81-87. doi: [10.1080/00050060500094605](https://doi.org/10.1080/00050060500094605)
- Voce, A., Calabria, B., Burns, R., Castle, D., & McKetin, R. (2019). A Systematic Review of the Symptom Profile and Course of Methamphetamine-Associated Psychosis. *Subst Use Misuse, 54*(4), 549-559. doi: [10.1080/10826084.2018.1521430](https://doi.org/10.1080/10826084.2018.1521430)